



Did you
KNOW

Cravings are one of the main reasons diets fail

Help control your cravings and
lose weight with **CONTRAVE[®]**

Actual patient. Results not typical. Across three studies, 46% of patients taking CONTRAVE with diet and exercise lost 5% or more of their body weight (vs 23% of patients on placebo).

CONTRAVE (naltrexone HCl/bupropion HCl) is a prescription weight-loss medicine that may help adults with obesity (BMI greater than or equal to 30 kg/m²), or who are overweight (BMI greater than or equal to 27 kg/m²) with at least one weight-related medical condition, lose weight and keep the weight off. CONTRAVE should be used along with diet and exercise.

Please see complete boxed warning for suicidal thoughts and behaviors and other Important Safety Information on page 7.

#1

PRESCRIBED
WEIGHT-LOSS BRAND*



Contrave[®]
(naltrexone HCl/bupropion HCl)
8 mg/90 mg • Extended-Release Tablets

*IQVIA NPA Branded Weight Loss Medicine Audit (Jan 2017 to July 2018).

Cravings and hunger are very different weight-loss challenges

Cravings ARE:

WANTING to eat specific foods. Your brain tells you it wants these specific foods even when you are not hungry

Cravings ARE NOT the same as:

Hunger, which is your brain and body telling you they need nutrition and any food will satisfy this **NEED**

You may experience cravings in different ways:



As a **reward** for something good

Let's get some ice cream to celebrate!



In certain **situations** or as part of a habit

Watching TV with a soda and a bag of potato chips.



When experiencing certain **emotions**


Chocolate calls my name when I'm stressed, bored, or lonely.



Do any of these sound familiar?

Ask your doctor today about a treatment option that may help you control your cravings and lose weight.

DID YOU KNOW...



DON'T WAIT!

Talk to your healthcare provider about ways to help control your cravings to lose weight

Share the times and situations you find yourself struggling with cravings.

For example, do you...

1

Feel like you deserve to eat a special treat when something good happens?

2

Want to snack whenever you sit down to watch TV?


3

Find yourself looking for comfort food when you're stressed, lonely, or bored?

Take 2 minutes to fill out the Discussion Guide in this booklet and review it with your healthcare provider.

If you're trying to lose weight and keep it off, 2 areas of your brain could be working against you*

- The hunger center (hypothalamus) can either reduce or trigger **hunger**



- The mesolimbic reward system can create **cravings** even when you're not hungry

*Other areas of the brain may be involved in weight loss.


Have you heard about set point theory?

Your body is thought to be programmed to defend against weight loss and maintain a certain range of body weight—a range that can be determined by your heaviest past weight.

This may be why hunger, and especially cravings, increase when dieting!


When weight loss occurs, the brain causes cravings that seem uncontrollable and increases hunger as a way to bring the body back to its predetermined set point.

Your set point may increase as you gain weight.



CONTRAVE is believed to work on 2 areas of your brain to reduce hunger and help control cravings*

The **hypothalamus (hunger center)** to reduce hunger



The **mesolimbic reward system** to help control cravings

*The exact neurochemical effects of CONTRAVE leading to weight loss are not fully understood.

2-4x
MORE WEIGHT LOST
along with diet and exercise

- Across 3 studies, patients who were overweight or struggling with obesity lost approximately **2-4x more** weight over one year by adding CONTRAVE than with diet and exercise alone
 - Nearly half of patients taking CONTRAVE lost 5% or more body weight and kept it off (vs 23% of patients taking placebo). Individual results may vary

MORE THAN
70%

of responders† with a BMI of 30.0-34.9 were able to move into a healthier, non-obese category (BMI 29.9 or less) after 1 year of CONTRAVE

†Patients who experienced ≥5% weight loss by Week 16.

Important Safety Information

Do not take CONTRAVE if you:

- have uncontrolled hypertension
- have or have had seizures or an eating disorder
- use other medicines that contain bupropion
- are dependent on opioid pain medicines, use medicines to help stop taking opioids, or are in opiate withdrawal
- drink a lot of alcohol and abruptly stop drinking, or take sedatives, benzodiazepines, or anti-seizure medicines and you abruptly stop using them
- are taking monoamine oxidase inhibitors (MAOIs). **Do not** start CONTRAVE until you have stopped taking your MAOI for at least 14 days
- are allergic to any of the ingredients in CONTRAVE
- are pregnant or planning to become pregnant or are breastfeeding

Before taking CONTRAVE, tell your healthcare provider about all of your current or past health conditions.

Please see additional Important Safety Information, including boxed warning, on page 7.


3 ways to connect with CONTRAVE

1

Visit us

at [CONTRAVE.com/health-and-fitness](https://www.contrave.com/health-and-fitness)

for diet and exercise tips, success stories from actual CONTRAVE patients, and more



2

Sign up

at [CONTRAVE.com/sign-up](https://www.contrave.com/sign-up)

for weight-loss tips and reminders to help you stay on track



3

Follow CONTRAVE


on Facebook and Instagram for tips to stay on track



CONTRAVE (naltrexone HCl/bupropion HCl) is a prescription weight-loss medicine that may help adults with obesity (BMI greater than or equal to 30 kg/m²), or who are overweight (BMI greater than or equal to 27 kg/m²) with at least one weight-related medical condition, lose weight and keep the weight off. CONTRAVE should be used along with diet and exercise.

Please see Important Safety Information and full Prescribing Information, including the boxed warning for suicidal thoughts and behaviors.


Contrave[®]
(naltrexone HCl/bupropion HCl)
8 mg/90 mg • Extended-Release Tablets



Are cravings keeping you from your weight-loss goals?

Check all that apply:

- I'm good all day, but then I reach for ice cream at night
- I often celebrate or reward myself with food
- When I am stressed or upset, I reach for comfort foods
- During holidays or special occasions, I find myself eating even when I'm not hungry

Other weight-related issues:

- My weight is holding me back from fully enjoying my life
 - I feel like my weight interferes with my ability to feel intimate with my partner
 - At work, I feel like people judge me for my weight, not for my abilities
 - My weight affects my self-confidence
 - Due to my weight, I feel like I can't keep up with my kids or friends as much as I'd like to
-



If you checked any of the above, cravings and/or weight-related struggles may be impacting your life. Talk to your doctor today about a treatment option that may help you control cravings and lose weight.

A HEALTHIER WEIGHT...

...gives you more to feel good about!

By working toward a healthier weight, you can help reduce your risk or symptoms of many weight-related conditions that can impact your health and everyday life, including:

Sleep apnea

Shortness of breath

Foot pain

Hip pain

Knee pain

Heartburn

Acid reflux

Joint pain

Your doctor will use BMI (body mass index) to assess your weight.

Ask your doctor how a lower BMI can be beneficial to your health and everyday life.

Study design for CONTRAVE clinical trials

Study 1 (COR-I): In this 56-week study, the group taking CONTRAVE along with diet and exercise lost 5.4% of their body weight (on average) compared with the placebo group who lost 1.3% (on average) with diet and exercise alone. Additionally, 42% of the CONTRAVE users lost at least 5% of their total body weight (while 17% of the placebo group lost at least 5% of their total body weight with diet and exercise). For participants who remained on CONTRAVE for the whole study, average weight loss was 8.1% or approximately 18 pounds, which was 4 times more weight than participants taking placebo.

Study 2 (COR-BMOD): In this 56-week study, all patients participated in an intensive diet and exercise program, including group visits. At 56 weeks, the CONTRAVE users lost 8.1% (on average) of their total body weight compared with a body weight loss of 4.9% (on average) for the placebo group. Additionally, 57% of those who took CONTRAVE lost at least 5% of their total body weight (while 43% of those on placebo lost at least 5% of their total body weight). For participants who remained on CONTRAVE for the whole study, average weight loss was 11.5% or 27 pounds.

Study 3 (COR-Diabetes): In this 56-week study of obese or overweight patients with type 2 diabetes, the group taking CONTRAVE along with diet and exercise lost 3.7% of their body weight (on average) compared with the placebo group who lost 1.7% (on average) with diet and exercise alone after 56 weeks. Additionally, 36% of the CONTRAVE users lost at least 5% of their total body weight (while 18% of the placebo group lost at least 5% of their total body weight). CONTRAVE users also had a reduction in HbA1c of 0.6% (compared with a reduction of 0.1% in the placebo group) at 56 weeks. CONTRAVE is not indicated to treat diabetes.

Important Safety Information

One of the ingredients in CONTRAVE, bupropion, may increase the risk of suicidal thinking in children, adolescents, and young adults. CONTRAVE patients should be monitored for suicidal thoughts and behaviors. In patients taking bupropion for smoking cessation, serious neuropsychiatric adverse events have been reported. **CONTRAVE is not approved for children under 18.**

Stop taking CONTRAVE and call your healthcare provider right away if you experience thoughts about suicide or dying; depression, or anxiety; panic attacks; trouble sleeping; irritability; aggression; mania; or other unusual changes in behavior or mood.

Do not take CONTRAVE if you:

- have uncontrolled hypertension
- have or have had seizures or an eating disorder
- use other medicines that contain bupropion
- are dependent on opioid pain medicines, use medicines to help stop taking opioids, or are in opiate withdrawal
- drink a lot of alcohol and abruptly stop drinking, or take sedatives, benzodiazepines, or anti-seizure medicines and you abruptly stop using them
- are taking monoamine oxidase inhibitors (MAOIs). **Do not** start CONTRAVE until you have stopped taking your MAOI for at least 14 days
- are allergic to any of the ingredients in CONTRAVE
- are pregnant or planning to become pregnant or are breastfeeding

Before taking CONTRAVE, tell your healthcare provider about all of your current or past health conditions.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. **Do not take any other medicines while you are taking CONTRAVE unless your healthcare provider says it is okay.**

If you have a seizure while taking CONTRAVE, stop taking CONTRAVE and call your healthcare provider right away.

Additional serious side effects may include: opioid overdose or sudden opioid withdrawal; severe allergic reactions; increases in blood pressure or heart rate; liver damage or hepatitis; manic episodes; visual problems (glaucoma); and increased risk of low blood sugar (hypoglycemia) in people with type 2 diabetes mellitus who take certain medicines to treat their diabetes.

The most common side effects of CONTRAVE include nausea, constipation, headache, vomiting, dizziness, trouble sleeping, dry mouth, and diarrhea.

These are not all the possible side effects of CONTRAVE. Please refer to the full Prescribing Information, including Medication Guide, for CONTRAVE or talk to your doctor.


You are encouraged to report negative side effects of drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

CONTRAVE IS THE #1 PRESCRIBED WEIGHT-LOSS BRAND*

Help control your cravings and lose weight with CONTRAVE

Want

Need



- Cravings and hunger are very different weight-loss challenges
 - Cravings are wanting specific foods regardless of whether you are hungry
 - Hunger is your body telling you it needs nutrition and any food will satisfy this need

- CONTRAVE is believed to work on 2 areas of your brain to reduce hunger and help control cravings
 - The hypothalamus (hunger center) to reduce hunger
 - The mesolimbic reward system to help control cravings

The exact neurochemical effects of CONTRAVE leading to weight loss are not fully understood.

2-4x
MORE WEIGHT LOST
along with diet and exercise

- Across 3 studies, patients who were overweight or struggling with obesity lost approximately 2-4x more weight over one year by adding CONTRAVE than with diet and exercise alone
 - Nearly half of patients taking CONTRAVE lost 5% or more body weight and kept it off (vs 23% of patients taking placebo). Individual results may vary

*IQVIA NPA Branded Weight Loss Medicine Audit (Jan 2017 to July 2018).

Don't wait.
Ask your doctor about CONTRAVE today!

Important Safety Information

Additional serious side effects may include: opioid overdose or sudden opioid withdrawal; severe allergic reactions; increases in blood pressure or heart rate; liver damage or hepatitis; manic episodes; visual problems (glaucoma); and increased risk of low blood sugar (hypoglycemia) in people with type 2 diabetes mellitus who take certain medicines to treat their diabetes.

Please see additional Important Safety Information, including boxed warning, on page 7.



Contrave[®]
(naltrexone HCl/bupropion HCl)
8 mg/90 mg • Extended-Release Tablets

**Nalprozion**
PHARMACEUTICALS

CONTRAVE[®] is a registered trademark of Nalprozion Pharmaceuticals, Inc.
All other trademarks are the property of their respective owners.
© 2018 Nalprozion Pharmaceuticals, Inc. CON-0245-006 12/2018 Printed in U.S.A.